



15th September 2015 - 14:00 h
CFEL – Building 99, seminar room I-III

Cedric Weber

King's College London, UK - Department of Physics

Dynamical mean field theory extended to nano-scopic systems and molecules

Phenomena that are connected to quantum mechanics, such as magnetism, transport, and the effect of impurity atoms and disorder, and their relation to material design and energy needs are important for almost every branch of the industry.

Density functional theory (DFT) was successful at making accurate predictions for many materials, in particular compounds which have a metallic behaviour. DFT combines high accuracy and moderate computational cost, but the computational effort of performing calculations with conventional DFT approaches is still non negligible and scales with the cube of the number of atoms.

A recent optimised implementation of DFT was however shown to scale linearly with the number of atoms (ONETEP), and opened the route to large scale DFT calculations for molecules and nanoscopic systems.

Nonetheless, one bottleneck of DFT and ONETEP, is that it fails at describing well some of the compounds where strong correlations are present, in particular because the computational scheme has to capture both the band-like character of the uncorrelated part of the compound and the Mott-like features emerging from the local strongly correlated centres. A recent progress has been made in this direction by the dynamical mean-field theory (DMFT), that allows to describe the two limits (metal and insulator) in a remarkable precise way when combined with DFT.

Recently, the TOSCAM package, a DMFT implementation for ONETEP, was tested and applied to metalloporphyrin systems. Extensions of DMFT to methods important for quantum chemistry analysis, such as the natural bond analysis, will be discussed.

References : PRL 110, 106402 ' 13 ; PRL 108, 256402 ' 12

Host: Dwayne Miller / CUI